

| Annex 1: data collection proforma | | | | | | | | | | | |
|---|---------|-----------------------|--|------------------------------|--------|------------------------|---------|------------------|-------------|--------------|--------------------|
| Variable | | | | | | | | | | | |
| History | | | | | | | | | | | |
| Gender | | | | | | M | F | | | | |
| Age (years) | | | | | | | | | | | |
| Hair loss | | | | | | Yes | No | | | | |
| Duration of hair loss | | | | | | | | | | | |
| Family history of hair loss.. | | | | | | Yes | No | | | | |
| Did hair loss start before or after SLE diagnosis | | | | | | Before | After | | | | |
| Are you still losing hair | | | | | | | | | | | |
| Any regrowth following treatment | | | | | | | | | | | |
| Any regrowth following treatment | | | | | | | | | | | |
| How long have you been on treatment? | | | | | | | | | | | |
| Clinical evaluation | | | | | | | | | | | |
| Normal hair | | | | | | Yes | No | | | | |
| Hair loss | | | | | | Yes | No | | | | |
| Pattern of hair loss | | | | | | Yes | No | | | | |
| Hair shaft changes | | | | | | | | | | | |
| Decreased follicular unit number (one hair in >20% of the follicles/visual field) | | | | | | Yes | No | | | | |
| Thin hair (decreased diameter in >20% of the follicles/visual field) | | | | | | Yes | No | | | | |
| Hypopigmented hair | | | | | | Yes | No | | | | |
| Hair loss type | | | | | | | | | | | |
| Patch <4 patches | | | | | | Yes | No | | | | |
| Mild diffuse (widening of central area) | | | | | | Yes | No | | | | |
| Severe diffuse (>50% of scalp) | | | | | | Yes | No | | | | |
| Skin | | | | | | | | | | | |
| Acute LE | | | | | | Yes | No | | | | |
| Subacute LE | | | | | | Yes | No | | | | |
| Chronic LE | | | | | | Yes | No | | | | |
| Lichenoid LE | | | | | | Yes | No | | | | |
| Mucosal LE (oral, nasal, conjunctival, genital) | | | | | | Yes | No | | | | |
| Chilblain LE | | | | | | Yes | No | | | | |
| Skin examination | | | | | | | | | | | |
| Malar rash | | Ulcers | | Blisters | | Discoid lesions | | Calcinosis cutis | | Oral ulcers | Fingertip erythema |
| Vasculitic rash | | Scaly patches/papules | | Scarring Alopecia | | Photosensitivity | | Lupus profundus | | Microthrombi | |
| Nailfold telangiectasias | | Raynaud's phenomenon | | Edema of the periungual skin | | Sausage shaped fingers | | | | | |
| Trichoscopy findings: | | | | | | | | | | | |
| Arborizing red loops | | | | | | | | | | | |
| Prominent (same diameter as terminal hair) | | | | Yes | | | No | | | | |
| Thin (<diameter of hair) | | | | Yes | | | No | | | | |
| Thick (>diameter of hair) | | | | Yes | | | No | | | | |
| Scalp pigmentary changes | | | | | | | | | | | |
| Normal/no hair loss area | | | | | | Hair loss area | | | | | |
| Area | Frontal | Rt temporal | | Lt temporal | Vertex | Occipital | Frontal | Rt temporal | Lt temporal | Vertex | Occipital |

| | | | | | | | | | | |
|--|--|--|--|--|-----|----|--|--|--|--|
| Black dots | | | | | | | | | | |
| White dots | | | | | | | | | | |
| Yellow dots | | | | | | | | | | |
| Large yellow dots (areas with absent follicles and scale | | | | | | | | | | |
| Brown scattered pigments | | | | | | | | | | |
| Brown peripilar sign | | | | | | | | | | |
| Follicular red dots | | | | | | | | | | |
| Blue-grey pigmentation | | | | | | | | | | |
| Peripilar cast | | | | | | | | | | |
| Short re-growing hair | | | | | Yes | No | | | | |
| Preserved honeycomb pigmentation | | | | | Yes | No | | | | |