

Images in clinical medicine



Point-of-care ultrasound detection of catheter-related tricuspid valve infective endocarditis in a patient undergoing haemodialysis

 Evans Nii Ayitey MacCready,  Isaac Nene Keteku

Corresponding author: Evans Nii Ayitey MacCready, Department of Internal Medicine, Cape Coast Teaching Hospital, Cape Coast, Ghana. maccvans@gmail.com

Received: 14 Jan 2026 - **Accepted:** 27 Jan 2026 - **Published:** 06 Feb 2026

Keywords: Infective endocarditis, catheter-related infection, point-of-care ultrasound

Funding: This work received no specific grant from any funding agency in the public, commercial, or non-profit sectors.

Copyright: Evans Nii Ayitey MacCready et al. Pan African Medical Journal (ISSN: 1937-8688). This is an Open Access article distributed under the terms of the Creative Commons Attribution International 4.0 License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Cite this article: Evans Nii Ayitey MacCready et al. Point-of-care ultrasound detection of catheter-related tricuspid valve infective endocarditis in a patient undergoing haemodialysis. Pan African Medical Journal. 2026;53(66). 10.11604/pamj.2026.53.66.51113

Available online at: <https://www.panafrican-med-journal.com//content/article/53/66/full>

Point-of-care ultrasound detection of catheter-related tricuspid valve infective endocarditis in a patient undergoing haemodialysis

Evans Nii Ayitey MacCready^{1&}, Isaac Nene Keteku¹

¹Department of Internal Medicine, Cape Coast Teaching Hospital, Cape Coast, Ghana

&Corresponding author

Evans Nii Ayitey MacCready, Department of Internal Medicine, Cape Coast Teaching Hospital, Cape Coast, Ghana

Image in medicine

A 32-year-old man with end-stage kidney disease secondary to chronic glomerulonephritis, on maintenance haemodialysis via a right internal jugular central venous catheter for three months, presented with a one-day history of worsening dyspnoea, fever, chills, non-productive cough, and central chest pain. On examination, he was febrile with a temperature of 38.6°C, tachycardic at 126 beats per minute, and his blood pressure was 150/98 mmHg. His oxygen saturation was 88% on

room air, which improved to 96% with supplemental oxygen. Chest examination findings were consistent with bilateral pleural effusions. Jugular venous pressure was not elevated, and no murmurs were heard. Point-of-care ultrasound demonstrated a large echogenic mass attached to the anterior leaflet of the tricuspid valve consistent with vegetation (red arrow), with the tip of the central venous catheter visualized within the right atrium (yellow arrow). Mild pericardial effusion and bilateral pleural effusions were also noted. Laboratory evaluation revealed marked leukocytosis with a white blood cell count of $28.6 \times 10^9/L$ and neutrophil predominance, with severe

anaemia (haemoglobin 7.2 g/dL). Blood cultures were obtained for suspected infective endocarditis following the ultrasound findings, and empiric intravenous antibiotic therapy was initiated promptly. However, the patient deteriorated rapidly and died the following day. This image highlights the importance of maintaining a high index of suspicion for catheter-related infective endocarditis in patients undergoing haemodialysis. It demonstrates the critical role of point-of-care ultrasound in the early detection of life-threatening cardiac complications associated with central venous catheters.




Figure 1: point-of-care ultrasound image demonstrating a tricuspid valve vegetation (red arrow) and the tip of a central venous catheter in the right atrium (yellow arrow)